DOI: 10.56871/UTJ.2024.20.52.005
УДК 616.831-005.4+616.98+616.936+616-056.5+612.39

НУТРИТИВНЫЙ СТАТУС ШКОЛЬНИКОВ СЕЛЬСКОЙ МЕСТНОСТИ ЧАРСАДДА, ПАКИСТАН

© Шакир Улла[1], Зишан Улла[1], Эйдул Ахад[2], Али Басит[3], Умайр Ислам[4], Мухаммад Ильяс[5], Усман Саид[6], Икбал Хан Рахман[7], Мухаммад Нур[8], Мухаммад Икбал[9], Хамза Хан[1]

[1] Абасинский университет. 25300, Пешаваре, Пакистан
[2] Сельскохозяйственный университет. 25000, Пешаваре, Пакистан
[3] Медицинский колледж. 25000, Спингар, Пакистан
[4] Сельскохозяйственный университет Синда. 70060, Тандо Джам, Пакистан
[5] Клиническая больница Хайбер. 25120, Пешаваре, Пакистан
[6] Международный университет Рифа. 23100, Малаканд, Пакистан
[7] Государственный колледж Университета. 54000, Лахор, Пакистан
[8] Университет Сват. 19220, Пакистан
[9] Кохатский университет науки и технологий. 26000, Кохат, Пакистан

Контактная информация: Shakir Ullah — врач. E-mail: shakirullah1992@gmail.com
ORCID: https://orcid.org/0000-0003-1731-9344

Для цитирования: Улла Ш., Улла З., Ахад Э., Басит А., Ислам У., Ильяс М., Саид У., Рахман И.Х., Нур М., Икбал М., Хан Х. Нутритивный статус школьников сельской местности Чарсадда, Пакистан // Университетский терапевтический вестник. 2024. Т. 6. № 3. С. 55–61. DOI: https://doi.org/10.56871/UTJ.2024.20.52.005

Поступила: 01.03.2024 ***Одобрена: 18.05.2024*** ***Принята к печати: 28.06.2024***

РЕЗЮМЕ. ***Введение.*** Малярия, анемия и недоедание являются актуальными проблемами здравоохранения, которые характеризуются высокой заболеваемостью и смертностью, большей распространенностью среди детей, особенно в Африке. ***Цель исследования*** — изучение распространенности и факторов риска развития малярии, анемии и недоедания среди детей, посещающих школу в Чарсадде (город Хайбер, Пакистан). ***Материалы и методы.*** В 2023 году в исследование было включено в общей сложности 400 детей в возрасте от 5 до 10 лет. Инфицирование малярией было подтверждено с помощью тестов экспресс-диагностики. Уровень гематокрита определяли с помощью центрифугирования, гемоглобин рассчитывали стандартным способом. Пищевой статус определяли с помощью антропометрических измерений, демографические характеристики оценивали с помощью анкетирования. Анемия и недостаточность питания определялись в соответствии со стандартами Всемирной организации здравоохранения. Для определения связи между прогностическими переменными и первичными исходами был использован логистический регрессионный анализ. ***Результаты.*** В ходе исследовательской работы было отобрано 400 детей для оценки малярии и состояния питания, малярийная инфекция была выявлена у 120 учащихся (12%). Наблюдалась высокая распространенность анемии — 105 (10,5%), мальнутриции — 90 (9%) и задержки роста — 70 (7%), в то время как истощение — 2 (0,2%) и дефицит массы тела — 20 (2%) встречались с очень низкой частотой. ***Выводы.*** Малярийная инфекция, анемия и недоедание наблюдались у 12, 10,5 и 9% детского населения соответственно. Насколько нам известно, это первое исследование, демонстрирующее состояние здоровья детей, посещающих школу в округе Чарсадда, а именно: распространенность малярии, анемии, недоедания и их сочетания у детей 5–10 лет.

КЛЮЧЕВЫЕ СЛОВА: анемия, малярия, недостаточное питание, задержка роста, недостаточный вес

NUTRITION STATUS OF SCHOOL-GOING CHILDREN IN THE RURAL AREA OF CHARSADDA, PAKISTAN

© *Shakir Ullah[1], Zeeshan Ullah[1], Eidul Ahad[2], Ali Basit[3], Umair Islam[4], Muhammad Ilyas[5], Usman Saeed[6], Iqbal Khan Rahman[7], Muhammad Noor[8], Muhammad Iqbal[9], Hamza Khan[1]*

[1] Abasyn University. Peshawar Pakistan 25300
[2] Agriculture University. Peshawar Pakistan 25000
[3] Medical College. Spinghar, Pakistan 25000
[4] Sindh Agriculture University. Tando Jam Pakistan 70060
[5] Khyber Teaching Hospital. Peshawar Pakistan 25120
[6] Riphah International University. Malakand Campus Pakistan 23100
[7] Government College University. Lahore Pakistan 54000
[8] University of Swat. Pakistan 19220
[9] Kohat University of Science & Technology. Kohat Pakistan 26000

Contact information: Shakir Ullah — Doctor. E-mail: shakirullah1992@gmail.com
ORCID: https://orcid.org/0000-0003-1731-9344

For citation: Ullah Sh., Ullah Z., Ahad E., Basit A., Islam U., Ilyas M., Saeed U., Rahman I.Kh., Noor M., Iqbal M., Khan H. Nutrition status of school-going children in the rural area of Charsadda, Pakistan. University Therapeutic Journal. 2024;6(3):55–61. DOI: https://doi.org/10.56871/UTJ.2024.20.52.005

Received: 01.03.2024 ***Revised: 18.05.2024*** ***Accepted: 28.06.2024***

ABSTRACT. ***Introduction.*** Malaria, anemia, and malnutrition are global health challenges with significant morbidity and mortality, and higher rates among children particularly in Africa. ***This study*** evaluated the prevalence and risk factors of malaria infection, anemia, and malnutrition among school-going children of Charsadda (a city of Khyber, Pakistan). ***Materials and methods.*** A total of 400 children of aged 5–10 years were included in the study in 2023. Malaria infection was confirmed by rapid diagnostic tests. Hematocrit level was obtained using a centrifuge microhematocrit, and converted to hemoglobin using standard conversion. Nutritional status was determined by the anthropometric measurements, and demographic characteristics were obtained by questionnaire. Anemia and malnutrition were defined according to the World Health Organization standards. Logistic regression analysis was used to determine association between the predictor variables and primary outcomes. Results and conclusion: In current research work 400 children were selected for the assessment of malaria and nutritional status, malaria infection was found more frequent among the students 120 (12%). In addition, high prevalence of anemia 105 (10.5%), malnutrition 90 (9%), and stunting 70 (7%) was observed, while wasting 2 (0.2%) and underweight 20 (2%) had very low frequency. Malaria infection, anemia, and malnutrition were observed in 12, 10.5 and 9% of the children population, respectively. To the best of our knowledge, this is the first study showing the current health problems among the school-going children of district Charsadda the prevalence of malaria, anemia and malnutrition and their co-existence in children 5–10 years.

KEYWORDS: anemia, malaria, malnutrition, stunting, underweight

INTRODUCTION

Rate of malaria infection declined from 22% (146 million individuals) in 2005 to 13% (114 million individuals) in 2015 in sub-Saharan Africa [1]. In Cameroon, malaria burden and transmission intensity are heterogeneous with spatial and temporal variations between altitudes and geographical areas, with varied prevalence rates from one area to another [2]. Like many sub-Saharan countries, prevalence of malaria has dropped in Cameroon by using insecticide-treated nets (ITN) in 2007 [3–6]. According to the follow-up study of Sumbele et al. [3], prevalence of malaria parasitemia dropped from 85.4% in 2006 to 36.6% in 2013 with a relative risk reduction of 57.2% in the Mount Cameroon area. Nevertheless, malaria still remains a major killer of children in this country and is estimated to take the life of a child every two minutes [1].

Malaria, anemia, and undernutrition are associated with morbidity and mortality, with higher rates among children, particularly in sub-Saharan Africa [7–9]. Anemia is defined as a condition in which the oxygen-carrying capacity of red blood cells is insufficient to meet the body's physiologic needs due to low blood hemoglobin concentration [7]. This condition affects individuals and has significant adverse health consequences, as well as adverse impacts on social and economic development [10].

Childhood anemia is considered a severe public health problem in Sub-Saharan Africa (62.5%) and in particular Cameroon, where a prevalence of 63.2% was reported in 2011 [1]. Malaria causes a substantial proportion of anemia in malaria-endemic settings [11–13]. Notwithstanding, updating the role of malaria parasitemia in anemia prevalence when coverage of insecticide-treated nets is above 75% in Cameroon [14], will help the National Malaria Control Programs to plan proper management strategies by considering the heterogeneities in different localities. However, association of anemia burden with malaria, relative to other causes such as malnutrition, and its variation across different altitudes of Cameroon has not been established. On the other hand, nutritional status is closely associated with immune response to the infections, is an important determinant of risk and prognosis of infectious diseases, and is directly influenced by the infections [15]. Nonetheless, association of malnutrition with malaria has been contradictory. Sumbele et al. [16] reported that malnutrition is associated with a higher risk of Plasmodium infection, and infectious episodes contribute to deterioration of nutritional status. In contrast, some studies found no association between nutrition and subsequent mortality from malaria [9, 17]. In agreement, malnutrition and Plasmodium falciparum malaria frequently coexist in Sahelian countries, and are accounted for a large part of under-five morbidity and mortality during their concomitant peak seasons [18]. Based on the report of the United Nations Children's Fund in 2013, 38% of children below the age of 5 years suffer from chronic malnutrition or stunting in sub-Saharan Africa. Unfortunately, malaria and undernutrition are two major causes of childhood mortality in the region [19]. Anemia has also been reported as a significant determinant of stunting [20], which is the main type of malnutrition in young children [21]. Stunting is associated with impaired cognitive development, reduced academic achievement, and decreased physical work capacity in adulthood, with a financial burden on societies [22]. While the global stunting prevalence fell from 39.6 to 23.8% between 1990 and 2014, scenario is quite different in Africa, where an increasing trend is observed. Nevertheless, prevalence of stunting fell from 49.9% [6] to 17.1% [21] in some localities in the Mount Cameroon area.

Impact of nutritional status on malaria may differ due to the heterogeneity of populations, species of the parasite, and other factors involved in the host and parasite relationship.

AIM OF THE STUDY

This study aimed at determining the prevalence and intensity of malaria parasitemia, anemia, and malnutrition as well as identifying the risk factors for these public health concerns among children living in Charsadda (a city of Khyber, Pakistan). Anemia and malnutrition were defined according to the World Health Organization standards. Logistic regression analysis was used to determine associations between the predictor variables and primary outcomes.

MATERIALS AND METHODS

This observational cross-sectional study was conducted in Charsadda city from April to September 2023. It was approved by ethical committee of the Takht Bhai Institute of Health and Management Sciences and the Takht Bhai THQ Hospital (Pakistan). Informed consent was obtained from the teachers and participants. Verbal consent was obtained from the parent/caregivers after explaining the purpose, risks, and benefits of the study. Sample size was determined by a single population formula. Accordingly, 400 students of aged 5–10 years were selected from the Govt schools. Demographic information together with weight and height of the children were obtained by a questionnaire (Table 1), and plotted to gender specific, growth charts. School Health Program was carried out on a systematic basis in the Rural Block. Ages of the children were determined using school records. For nutritional status assessment, weight was measured by a floor type weighing scale, height was taken using a measuring tape applied to the wall from their back of heels to the head touching the wall. Malaria infection was confirmed by rapid diagnostic tests. The hematocrit level was obtained using a centrifuge microhematocrit and converted to hemoglobin using standard conversion. Nutritional status was determined from the anthropometric measurements collected. Food

Table 1

Indicators and cutoffs for assessment of malnutrition in the school-going children

Demographic information	**General information**	**Dietary Habits**
Name:	Do you have breakfast every day before going to school? (Yes/No)	What is your favorite healthy food?
Age:		
Gender:		
Grade / Class:	How many meals do you typically eat a day?	How often do you drink water a day?
School Name:		Less than 3 glasses
Anthropometric Measurements	Do you eat fruits and vegetables daily? (Yes/No)	3–5 glasses
Height (cm):		6–8 glasses
Weight (kg):	How often do you consume fast food? (Yes/No)	More than 8 glasses
BMI (Body Mass Index):		Are you aware of the importance of a balanced diet? (Yes/No)
		Do you receive any nutrition education at school? (Yes/No)

frequency questionnaire for Nutrition status was ailo used (Table 3). Anemia and malnutrition were defined according to the World Health Organization standards.

RESULTS AND DISCUSSION

In current research work 400 children were selected for the assessment of malaria and nutritional status, malaria infection was found more frequent among the students 120 (12%). In addition, high prevalence of anemia 105 (10.5%), malnutrition 90 (9%), and stunting 70 (7%) was observed, while wasting 2 (0.2%) and underweight 20 (2%) had very low frequency. Malaria infection, anemia, and malnutrition were observed in 12, 10.5 and 9% of the children population, respectively (Figure 1).

In age wise incidence of Malaria the lower age group were more infect by Malaria as compare to higher age group students 6.5 and 5.5% respectively (Table 2). While gender wise incidence of Malaria shows that the more boys were infected by Malaria as compare to girls, boys 70 (7%) and girls 50 (5%) (Table 2).

Overall assessment of nutritional status was found more frequent, malnutrition 90(9%), and stunting 70 (7%) was observed, while wasting 2 (0.2%) and underweight 20 (2%) had very low frequency. Malnutrition were observed in 12, 10.5 and 9% of the school-going children respectively (Figure 2).

In the current research work a total of 400 samples were isolated from school-going children. The samples collected from male and females students of age 5–10 years. Among the all samples malarial infection were found more frequent (120), leading by anemia (105), similarly malnutrition also with high prevalence rate (90) while wasting and underweight with very low




Fig. 1. Clinical characteristics of the study population

Table 2

Malaria infection prevalence with respect to sex and age

Parameter	Number	Prevalence (%)
Age		
0–5	65	6.5%
6–10	55	5.5%
Sex		
Girls	50	5%
Boys	70	7%

frequency of (2) and (20) respectively, stunting were also having frequency of (70). Out of 1000 children the number of male was slightly higher than females.

More males were malnourished (43.6%), underweight (18.1%) and stunted (42.6%) when compared with females 39.7, 7.1 and 37.2% respectively though the difference was not significant in malnutrition and stunting. The difference in prevalence of underweight among the sexes was significant (p=0.025) with the males having




Fig. 2. Over all frequency of malnutrition, wasting, underweight and stunting

Table 3

Food frequency questionnaire for Nutrition status of school-going children in the rural area of Charsadda, Pakistan

Food frequency questionnaire (ffq)				
Meal timings				
Do you take	Always	Sometimes	Rarely	Never
Pre-Breakfast	50	50	140	160
Breakfast	30	70	100	200
Brunch	10	30	60	300
Lunch	200	160	30	10
Tea time	150	100	110	40
Dinner	220	80	60	40
Post Dinner	50	100	150	100
Food response and reactions				
Cravings		Aversions		Unsuitability
300		60		40
Water intake (1 Glass = 250 ml)				

a higher prevalence. Bivariate analysis revealed children of the 0–5 years age group ($p < 0.001$) were significantly at odds of being malnourished. Anemia has also been testified as a important cause of stunting [20], which is the main type of malnutrition in young children [21]. Stunting is associated with impaired cognitive development, reduced academic achievement, and decreased physical work capacity in adulthood, with a negative cost on the economic development of societies [22]. While the global stunting prevalence fell from 39.6 to 23.8% between 1990 and 2014, the scenario is quite different in Africa, with an increase. Nevertheless, in some localities in the Mount Cameroon area, the prevalence of stunting fell from 49.9 to 17.1% [21]. The impact of nutritional status on malaria may differ due to the heterogeneity of the population under study, species of the parasite, and other factors involved in the host and parasite relationship. The study aimed at defining the prevalence and intensity of malaria parasitaemia, anaemia and malnutrition as well as identifying the risk factors for these public health concerns among children living in low versus high altitude settings in the Mount Cameroon area.

CONCLUSION

The available data indicate that malnutrition is a public health problem among the school-going children of the local area of District Charsadda, Pakistan. However, the available data, on micronutrient status, are limited. These findings highlight the need for nutrition involvements in school-going children and more high-quality research to assess nutritional status in this age group.

Anemia and malnutrition control should be integrated with existing malaria control and should include children above five years of age. In the current research work among the all malarial infection were found more frequent (120), leading by Anemia (105), similarly malnutrition also with high prevalence rate(90) while wasting and underweight with very low frequency of (1) and (15) respectively, stunting were also having frequency of (65). Out of 400 children the number of male was slightly higher than females.

To the best of our knowledge, this is the first study showing the current health problems among the school-going children of District Charsadda describing the prevalence of malaria, anemia and malnutrition and their co-existence in children 10 years and below. The key observation is that there is a limited study on the health problems of this population in District Charsadda. This research work however provides data for further studies and advance on public health among school-going children.

Благодарности

Мы ценим поддержку Департамента патологии THQ Тахт бахи Мардан в успехе исследования. Мы также хотели бы поблагодарить сотрудников Школы и руководителей школ за техническую поддержку.

Конфликт интересов

Авторы заявляют, что у них нет конфликта интересов.

Acknowledgements

We appreciate the support of the Department of Pathology THQ Takht bahi Mardan towards the success of the study. We will also like to thank the staffs of the School teachers and school heads for their technical support.

Competing interests

The authors declare that they have no competing interests.

ЛИТЕРАТУРА / REFERENCES

1. Apinjoh T.O., Anchang-Kimbi J.K., Mugri R.N., Tangoh D.A., Nyingchu R.V., Chi H.F. et al. The efect of insecticide treated nets (ITNs) on Plasmodium falciparum infection in rural and semi urban communities in the South West Region of Cameroon. PLoS One. 2015;10:e0116300.
2. Akiyama T., Pongvongsa T., Phrommala S., Taniguchi T., Inamine Y., Takeuchi R. et al. Asymptomatic malaria, growth status, and anaemia among children in Lao People's Democratic Republic: a cross-sectional study. Malaria Journal. 2016;15:499. DOI: 10.1186/s12936-016-1548-3.
3. Best C., Neufingerl N., Van Geel L., van den Briel T., Osendarp S. The nutritional status of school-aged children: why should we care? Food and nutrition bulletin. 2010;31(3):400–417. DOI: 10.1177/156482651003100303.
4. Boussougou-Sambe S.T., Awono-Ambene P., Tasse G.C., Etang J., Binyang J.A., Nouage L.D. et al. Physical integrity and residual bio-efcacy of used LLINs in three cities of the South-West Region of Cameroon 4 years after the first national mass-distribution campaign. Malaria Journal. 2017;16:31. DOI: 10.1186/s12936-017-1690-6.
5. Caulfeld L.E., Richard S.A., Black R.E. Undernutrition as an underlying cause of malaria morbidity and mortality in children less than five years old. The American journal of tropical medicine and hygiene. 2004;71(suppl 2):55–63. DOI: 10.1186/s12936-018-2492-1.
6. Ebai C.B., Kimbi H.K., Sumbele IUN., Yunga J.E., Lehman L.G. Epidemiology of Plasmodium falciparum malaria in the Ikata-Likoko area of Mount Cameroon: a cross sectional study. International Journal of Tropical disease & Health. 2016;16:1–12. DOI: 10.9734/IJTDH/2016/25890.
7. Eyong E.E.J., Kengne-Ouafo A.J., Chounna P.W., Datchoua-Poutcheu F.R., Wanji S. Altitudinal variation in the parasitological and entomological indices of malaria around Mount Cameroon, South West Region of Cameroon. Journal of Parasitology and Vector Biology 2016;8:74–85. DOI: 10.5897/JPVB2016.0242.
8. Kateera F., Ingabire C.M., Hakizimana E., Kalinda P., Mens P.F., Grobusch M.P. et al. Malaria, anaemia and under-nutrition: three frequently coexisting conditions among preschool children in rural Rwanda. Malarial Journal. 2015;14:440. DOI: 10.1186/s12936-015-0973-z.
9. Kimbi H.K., Nkesa S.B., Ndamukong-Nyanga J.L., Sumbele IUN., Atashali J., Atanga M.B. Knowledge and perceptions towards malaria prevention among vulnerable groups in the Buea Health District, Cameroon. BMC Public Health. 2014;14:883. DOI: 10.1186/1471-2458-14-883.
10. Martorell R., Rivera J., Kaplowitz H., Pollitt E. Long-term consequence of growth retardation during early

childhood. In: Hernandez M., Argente J., editors. Human growth: basic and clinical aspects. Amsterdam: Elsevier Science Publ; 1992:143–9.
11. Nkuo-Akenji T.K., Sumbele I., Mankah E., Njunda A., Samje M., Kamga L. The burden of malaria and malnutrition among children less than 14 years of age in a rural village of Cameroon. African Journal of Food, Agriculture, Nutrition and Development. 2008;8:252–64. DOI: 10.4314/ajfand.v8i3.19110.
12. Nyaaba D.A., Brown P.K., Agbenorhevi J.K. Assessing the association between nutritional status and asymptomatic malaria parasitaemia of children under five years in Ghana. Pyrex J Nutrition Metabolism. 2017;2:1–14. DOI: 10.1186/s12936-018-2492-1.
13. Sumbele I.U., Ning T.R., Bopda O.S., Nkuo-Akenji T. Variation in malariometric and red cell indices in children in the Mount Cameroon area following enhanced malaria control measures: evidence from a repeated crosssectional study. Malaria Journal. 2014;13:334. DOI: 10.1186/1475-2875-13-334.
14. Stevens G.A., Finucane M.M., De-Regil L.M., Paciorek C.J., Flaxman S.R., Branca F. et al. Global, regional, and national trends in haemoglobin concentration and prevalence of total and severe anaemia in children and pregnant and non-pregnant women for 1995–2011: a systematic analysis of population-representative data. Lancet Global Health. 2013;1:e16–25. DOI: 10.1016/S2214-109X(13)70001-9.
15. Sumbele IUN., Sama S.O., Kimbi H.K., Taiwe G.S. Malaria, moderate to severe anaemia, and malarial anaemia in children at presentation to hospital in the Mount Cameroon area: across-sectional study. Anemia.2016;2016:5725634. DOI: 10.1155/2016/5725634.
16. Sumbele IUN., Kimbi H.K., Ndamukong-Nyanga J.L., Nweboh M., Anchang-Kimbi J.K., Lum E. et al. Malarial anaemia and anaemia severity in apparently healthy primary school children in urban and rural settings in the Mount Cameroon area: cross sectional survey. PLoS One. 2015;10:e0123549. DOI: 10.1371/journal.pone.0123549.
17. UNICEF. Annual Report 2015. New York: 2015. https://www.unicef.org/ publications/index_92018.html. Accessed 27 Apr 2024. DOI: 10.1186/s12936-018-2492-1.
18. Unicef. Improving child nutrition. The achievable imperative for global progress. 2013. New York: UNICEF; 2016. https://www.unicef.org/nutrition/ index_68661.html. Accessed 27 Apr 2024.
19. UNICEF. United Nations interagency group for child mortality estimation. Levels and trends in child mortality. In. New York: United Nations Children's Fund; 2014. http://www.childmortality.org/fles_v21/download/ unicef-2013-child-mortality-report-LR 10_31_14_195. DOI: 20.500.12592/k9c67c.
20. WHO. World malaria report 2015. Geneva: World Health Organization; 2016. http://www.who.int/maaria/publications/world-malaria report2016/report/en/. Accessed 27 Apr 2024. DOI: 10.1186/s12936-016-1548-3.
21. Wanzira H., Katamba H., Okullo A.E., Agaba B., Kasule M., Rubahika D. Factors associated with malaria parasitaemia among children under 5 years in Uganda: a secondary data analysis of the 2014 Malaria Indicator Survey dataset. Malarial Journal. 2017;16:191. DOI: 10.1186/s12936-017-1847-3.
22. WHO. World Malaria Report. Geneva: World Health Organization; 2014. http://www.who.int/malaria/publications/world_malaria_report_2014/en/. Accessed 27 Apr 2024. DOI: 10.1186/s12936-018-2492-1.